

STREPTOCOCCAL SORE THROAT/SCARLET FEVER

Streptococcal sore throat (strep throat) and scarlet fever (a strep throat with a rash) are common infections in young children. These illnesses are usually not serious. However, complications such as rheumatic fever or kidney disease may rarely develop if children do not receive proper antibiotic treatment.

CAUSE: *Streptococcus* bacteria (Group A beta hemolytic strep)

SYMPTOMS: Sudden onset of fever, sore throat, swollen glands, headache, and abdominal pain. Nausea and vomiting may occur with severe cases. With scarlet fever, a very fine, bright red, raised rough texture rash (feels like sandpaper) is present. A fuzzy white tongue followed by a beefy-red 'strawberry tongue' may occur. The rash appears most often on the neck, chest, inner thigh, and in folds of the armpit, elbow, and groin. Later on, there may be peeling of the skin on the fingertips and toes.

SPREAD: Person-to-person from nose and throat secretions of infected persons (those with and without symptoms).

INCUBATION: It usually takes 1 to 3 days from the time a person is exposed until symptoms develop.

CONTAGIOUS PERIOD: Until at least a full 24 hours after treatment begins and fever is gone.



Until at least a full 24 hours after treatment begins and until child is without fever for 24 hours.



Prevention Control

1. If your child does not appear well or develops a sore throat and other symptoms listed above, keep him/her home and call your physician.
2. **Diagnosis:** Confirmed by identification of strep in the throat, either by throat culture, or by using a rapid test which can provide results the same day.
3. **Treatment:** If the culture is positive, a penicillin shot or antibiotics taken by mouth (usually 10 days) are prescribed. This treatment will help to prevent more serious illness such as rheumatic fever, which can damage the heart valves.

STREPTOCOCCAL SORE THROAT/SCARLET FEVER